

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich							
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM	
1,25-(OH)-Vitamin-D3	Serum		LIA	75	7	1	-	Sero	-	-	-	-	-	-
10-OH-Carbazepin(MHD)	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
17-OH-Progesteron i.S.	Serum		ELISA	>1			-	Sero	-	-	-	-	-	-
17β-Ostradiol (E2)	Serum		ECL	180	2	1	-	KC	Basis	-	-	-	-	-
1-OH-Alprazolam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
1-OH-Midazolam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
1-OH-Triazolam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
25-(OH)-Vitamin D	Serum		HPLC	28	4	2	-	-	-	HPLC	-	-	-	-
3α-Androstadiol-Glucuronid	Serum		ELISA	>1	1	-	-	Sero	-	-	-	-	-	-
3-OH-Bromazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
7-Amino-Clonazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
7-Aminoflunitrazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
7-Aminonitrazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
9-Hydroxy-Risperidon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
a1-Antitrypsin	Serum		NEPH	90	90	7	-	KC	-	-	-	-	-	-
a1-Fetoprotein (AFP)	Serum		ECL	180	14	5	-	KC	Basis	KC	-	-	-	-
a1-Mikroglobulin i.H.	Urin		NEPH	-	7	-	-	-	Liq	-	-	-	-	-
AAK gg Amphiphysin	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg Aquaporin-4 IgG i.S.°	Serum		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg Aquaporin-4 IgG im L.c.°	Liquor		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg CASPR2°	Serum		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg CV2 (CRMP5)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg cycl. citr. Peptide (CCP)	Serum		CLIA	>7	7	8h	-	Sero	-	-	-	-	-	-
AAK gg DPPX IgG im L.c.°	Liquor		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg DPPX-IgG°	Serum		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg ds-DNA	Serum		ELISA	-	14	7	-	Sero	Spez	-	-	-	-	-
AAK gg Endomysium IgA	Serum		IFT	>14	14	-	-	-	-	Viro	-	-	-	-
AAK gg Endomysium IgG	Serum		IFT	>14	14	-	-	-	-	Viro	-	-	-	-
AAK gg GAD (GAD65)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg GAD (GAD65)°	Serum		ELISA	>3	3	1	-	-	-	Viro	-	-	-	-
AAK gg Ganglioside: GM1, GM2, GM3, GM4, GD1a, (Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg glatte Muskulatur (ASMA)	Serum		IFT	>14	14	1	-	-	-	Viro	-	-	-	-
AAK gg glom. Basalmembran	Serum		EIA	>5	5	-	-	Sero	-	-	-	-	-	-
AAK gg GP 210 IgG	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-	-
AAK gg Insulin°	Serum		ELISA	>3	3	1	-	-	-	Viro	-	-	-	-
AAK gg Intrinsic Faktor°	Serum		IB	>5	5	1	-	Sero	-	-	-	-	-	-
AAK gg Jo-1	Serum		IB	>14	14	-	-	-	-	Viro	-	-	-	-
AAK gg Ku	Serum		IB	>14	14	-	-	-	-	Viro	-	-	-	-
AAK gg Leber/Pankreas (SLA / LP, IgG)	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-	-
AAK gg Lebercytosol (LC1)	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-	-
AAK gg Leberspez. Protein (LSP)	Serum		IFT	>14	14	1	-	-	-	Viro	-	-	-	-
AAK gg Leberzellmembran (LMA)	Serum		IFT	>14	14	1	-	-	-	Viro	-	-	-	-
AAK gg Liver-Kidney-Mikrosomen (LKM)	Serum		IFT	>14	14	1	-	-	-	Viro	-	-	-	-
AAK gg Liver-Kidney-Mikrosomen (LKM1)	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-	-
AAK gg Mitochondrien (AMA)	Serum		IFT	>14	14	1	-	-	-	Viro	-	-	-	-
AAK gg Mitochondrien Typ M2	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-	-
AAK gg MOG i.S.°	Serum		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg MOG im L.c.°	Liquor		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg mutiertes citrulliniertes Vimentin	Serum		ELISA	>5	5	-	-	Sero	-	-	-	-	-	-
AAK gg Neuronenkerne (Hu)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg Neuronenkerne (Ri)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg NMDA-NR1-Rezeptoren im L.c.°	Liquor		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg NMDA-NR1-Rezeptoren°	Serum		IFT	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg Pankreas-Inselzellen (ICA)	Serum		IFT	>14	14	1	-	-	-	Viro	-	-	-	-
AAK gg PNMA2 (Ma-2/Ta)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg Purkinjezellen (Tr)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg Purkinjezellen (Yo)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg Recoverin	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg SOX1	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg SP100	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-	-
AAK gg Thyreoglobulin (TAK)	Serum		CLIA	90	1	1	-	-	Spez	-	-	-	-	-
AAK gg Titin (MGT-30)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg TPO (MAK)	Serum		ECL	2a	8	8	ANA	KC	Basis	KC	-	-	-	-
AAK gg TSH-Rezeptor (TRAK)	Serum		ECL	360	6	7h	ANA	KC	Basis	KC	-	-	-	-
AAK gg Tyrosinphosphatase (IA-2)°	Serum		ELISA	>3	3	1	-	-	-	Viro	-	-	-	-
AAK gg Zink-Finger-4-Protein (ZIC-4)	Serum		IB	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg. AMPA-1/2-Rezeptoren im L.c.°	Liquor		IFT	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg. AMPA1-Rezeptoren i.S.*	Serum		IFT	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg. Annexin V (IgG)	Serum		EIA	>5	5	-	-	Sero	-	-	-	-	-	-
AAK gg. Annexin V (IgM)	Serum		EIA	>5	5	-	-	Sero	-	-	-	-	-	-
AAK gg. CASPR2 im L.c.°	Liquor		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg. GABA-B1/2-Rezeptoren°	Serum		IFT	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg. GABA-B1/B2-Rezeptoren im L.c.°	Liquor		IFT	90	14	-	-	-	Spez	-	-	-	-	-
AAK gg. Lgi1 im L.c.°	Liquor		IFT	90	14	-	-	-	Liq	-	-	-	-	-
AAK gg. LGI-1°	Serum		IFT	90	14	-	-	-	Liq	-	-	-	-	-
ACE i.S.	Serum		PHOT	180	30	4h	-	Sero	-	-	-	-	-	-
Adenoviren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-
Adenovirus 40/41 (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-
Agomelatin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-
aktueller PH-Wert	Kapillarblut		AMPE	-	-	10'	-	-	-	Häma	FI	HZ	-	-

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich						
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM
ALAT (GPT)	Serum		PHOT	7	7	3	ANA	KC	Basis	KC	FI	HZ	SM
Albumin	Serum		TURB	120	150	70	ANA	KC	Basis	KC	FI	-	-
Albumin (%)	Serum		ELPHO	60	10	3	-	Sero	Basis	-	-	-	-
Albumin i.H.	Urin		TURB	180	30	7	-	KC	Basis	KC	-	-	-
Albumin im L.c.	Liquor		TURB	180	3	-	-	-	Liq	-	-	-	-
Aldosteron i.S., basal	Serum		CLIA	28	5	1	-	-	Spez	-	-	-	-
Alkalische Phosphatase	Serum		PHOT	60	7	7	ANA	KC	Basis	KC	FI	HZ	SM
Alkohol (Ethanol) i.S.	Serum		PHOT	28	14	2	-	-	-	KC	FI	HZ	-
Alpha1-Globulin (%)	Serum		ELPHO	60	10	3	-	Sero	Basis	-	-	-	-
Alpha2-Globulin (%)	Serum		ELPHO	60	10	3	-	Sero	Basis	-	-	-	-
Alprazolam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Aluminium°	Sonstiges		AAS	-	90	-	-	AAS	-	-	-	-	-
Amiodaron	Serum		HPLC	28	4	-	-	-	-	HPLC	-	-	-
Amisulprid	Serum		LC/MS	90	7	3	-	-	-	HPLC	-	-	-
Ammoniak i. Pl.	EDTA	stationär: 30 Minuten	PHOT	-	-	30'	-	KC	-	KC	FI	HZ	-
Amphetamine i.H. (immunol.)	Urin		KIMS	>5	5	1	-	-	-	KC	-	-	-
Amphetamine i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-
Amyloid Aβ40 im L.c.	Liquor		CMIA	60	7	-	-	-	Liq	-	-	-	-
Amyloid Aβ42 im L.c.	Liquor		CMIA	60	7	-	-	-	Liq	-	-	-	-
ANA-Blot: Anti-PM/ScI	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: ds-DNA	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: Histone	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: Jo-1 (Anti-His-tRNA-Synthetase)	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: Ku	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: Mi-2	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: Nukleosomen	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: Scl-70 (Anti-Topoisomerase I)	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: SM-D1	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: SS-A/ Ro 52 kD	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: SS-A/ Ro 60 kD	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: SS-B/ La	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: U1-sn-RNP	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
ANA-Blot: Zentromere	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
Androstendion	Serum		ELISA	12m	5	-	-	Sero	-	-	-	-	-
Anti-C1q	Serum		EIA	>5	5	-	-	Sero	-	-	-	-	-
Anti-DFS70	Serum		IFT	>3	3	1	-	-	-	Viro	-	-	-
Anti-DGP IgA (Gliadin - AK IgA)	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-
Anti-DGP IgG (Gliadin - AK IgG)	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-
Anti-dsDNA	Serum		IB	>2	2	8h	-	-	-	Viro	-	-	-
Anti-HAV IgM	Serum		ECL	90	17	7	-	KC	Basis	KC	-	-	-
Anti-HAV-IgG+IgM	Serum		ECL	90	14	6	-	KC	Basis	KC	-	-	-
Anti-HBc	Serum		ECLIA	90	14	7	-	KC	Basis	KC	-	-	-
Anti-HBc-IgM	Serum		ECL	90	14	7	-	KC	Basis	KC	-	-	-
Anti-HBe	Serum		ECLIA	90	14	7	-	-	Basis	-	-	-	-
Anti-HBs-Titer	Serum		ECLIA	90	6	3	-	KC	Basis	KC	-	-	-
Anti-HIV-1/2+p24-Ag	Serum		EIA	90	28	7	-	KC	Basis	KC	-	-	-
Antikörpersuchtest	Vollblut		Aggl	360	10	1	-	-	-	Häma	FI	HZ	-
Anti-Müller-Hormon	Serum		ELISA	>6	6	1	-	KC	-	-	-	-	-
Antinukleäre AK (ANA)	Serum		IFT	>3	3	1	-	-	-	Viro	-	-	-
Antistreptolysin (ASL)	Serum		TURB	180	8	2	-	KC	Basis	KC	-	-	-
Antithrombin	Citrat		CHRO	2a	-	4h	-	KC	-	Häma	-	HZ	-
APC-Sensitivität (-Ratio)	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Apolipoprotein A1	Serum		TURB	56	8	1	-	KC	-	-	-	-	-
Apolipoprotein B	Serum		TURB	60	8	1	-	KC	-	-	-	-	-
Aripiprazol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
ASAT (GOT)	Serum		PHOT	90	7	4	ANA	KC	Basis	KC	FI	HZ	SM
ASCA	Serum		IB	>5	5	1	-	Sero	-	-	-	-	-
Astrovirus (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-
Atomoxetin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Barbiturate i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-
Bartonella henselae-IgG-AK°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-
Bartonella henselae-IgM-AK°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-
Basophile, absolut	EDTA		WhF	-	2	2	-	Häma	Basis	Häma	FI	HZ	-
Benzodiazepine i.H.(immunol.)	Urin		KIMS	>5	5	1	-	-	-	KC	-	-	-
Benzodiazepine i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-
Beta-1-Globulin (%)	Serum		ELPHO	60	10	3	-	Sero	Basis	-	-	-	-
Beta-2-Globulin (%)	Serum		ELPHO	60	10	3	-	Sero	Basis	-	-	-	-
Beta-CrossLaps (β-CTX)	EDTA-P		ECL	90	8h	6h	-	-	Basis	-	-	-	-
Bilirubin direkt	Serum		PHOT	180	7	2	ANA	KC	Basis	KC	FI	HZ	-
Bilirubin gesamt	Serum		PHOT	180	7	1	ANA	KC	Basis	KC	FI	HZ	SM
Bilirubin i.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM
Blastocystis hominis (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-
Blutgruppe	Vollblut		Aggl	-	10	1	-	-	-	Häma	FI	HZ	-
Blutsenkung (1h)	EDTA		PHOT	-	2	6h	-	KC	Basis	Häma	-	-	-
B-Lymphozyten (CD19)	EDTA		FC	-	-	1	-	-	Spez	-	-	-	-
Bocaviren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-
Bordetella parapertussis-DNA (PCR)	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-
Bordetella pertussis Toxin-IgA AK	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-
Bordetella pertussis Toxin-IgG AK	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-
Bordetella pertussis-DNA (PCR)	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich							
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM	
Borrelie Ak IgG	Serum		ELISA	-	14	7	-	-	Spez	-	-	-	-	-
Borrelie Ak IgM	Serum		ELISA	-	14	7	-	-	Spez	-	-	-	-	-
Borrelie Blot IgG	Serum		IB	60	14	7	-	-	Spez	-	-	-	-	-
Borrelie Blot IgM	Serum		IB	60	14	7	-	-	Spez	-	-	-	-	-
Borrelie-IgG i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-
Borrelie-IgG-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-
Borrelie-IgM i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-
Borrelie-IgM-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-
Brivaracetam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Bromazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Brucellen IgG Ak°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-	-
Brucellen IgM Ak°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-	-
Buprenorphin (qualitativ)	Urin		IA	>5	5	1	-	Sero	-	-	-	-	-	-
Bupropion	Serum		LC-MS/MS	90	1	5h	-	-	-	HPLC	-	-	-	-
c - ANCA (IFT)	Serum		IFT	>14	14	-	-	-	-	Viro	-	-	-	-
C. difficile Toxin-Gen (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-
C3-Komplement	Serum		RID	-	7	-	-	-	Liq	-	-	-	-	-
C4-Komplement	Serum		RID	-	7	-	-	-	Liq	-	-	-	-	-
CA 125	Serum		ECL	168	5	8h	-	KC	Basis	KC	-	-	-	-
CA 15-3	Serum		ECL	90	5	2	-	KC	Basis	KC	-	-	-	-
CA 19-9	Serum		ECL	90	14	5	-	KC	Basis	KC	-	-	-	-
CA 72-4	Serum		ECL	90	30	1	-	KC	-	-	-	-	-	-
Calcitonin	Serum		ECL	2a	1	4h	-	KC	-	-	-	-	-	-
Calcium i.H.	Urin		PHOT	21	4	2	-	-	Basis	KC	-	-	-	-
Calcium i.S.	Serum		PHOT	240	21	7	ANA	KC	Basis	KC	FI	HZ	SM	-
Calcium ionis.	Serum		AMPE	-	10	3	-	KC	-	Häma	-	-	-	-
Calprotectin i. Stuhl	Stuhl		CLIA	7	3	1	-	-	Mibi	-	-	-	-	-
Campylobacter (jejuni/coli) IgA	Serum		ELISA	>14	14	<1	-	-	-	Viro	-	-	-	-
Campylobacter (jejuni/coli) IgG	Serum		ELISA	>14	14	<1	-	-	-	Viro	-	-	-	-
Campylobacter spp. (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-
C-ANCA (PR3)	Serum		ELISA	-	14	7	-	Sero	Spez	-	-	-	-	-
Cannabinoide i.H. (immunol.)	Urin		KIMS	>5	5	1	-	-	-	KC	-	-	-	-
Cannabinoide i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-	-
Carbamazepin	Serum		LC-MS/MS	>2	2	8h	-	-	-	HPLC	-	-	-	-
Carbamazepin-Diol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Carbamazepinepoxyd	Serum		LC/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Carbohydrate deficient transferrin (CDT)	Serum		HPLC	28	7	1	-	-	-	HPLC	-	-	-	-
Cardiolipin - IgG	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-	-
Cardiolipin - IgM	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-	-
Cardiolipin-Mikrofl.-Test (VDRL)	Serum		IHA	-	7	-	-	-	Liq	-	-	-	-	-
CEA	Serum		ECL	180	14	7	-	KC	Basis	KC	-	-	-	-
Chlam. pneum. IgA Ak	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-	-
Chlam. pneum. IgG Ak	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-	-
Chlam. trach. IgA Ak	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-	-
Chlam. trach. IgG Ak	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-	-
Chlamydia pneumoniae Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-
Chlamydia trachomatis Multiplex	Abstrich		PCR	-	7	2	-	-	Mibi	-	-	-	-	-
Chlamydia trachomatis rRNA	Abstrich		NAT	-	-	30	-	-	PCR	-	-	-	-	-
Chlamydia trachomatis rRNA	Urin		NAT	-	-	30	-	-	PCR	-	-	-	-	-
Chlordiazepoxid	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Chlorid i.H./Konz.	Urin		ISE	>7	>7	7	-	-	Basis	KC	-	-	-	-
Chlorid i.S.	Serum		ISE	>7	7	7	ANA	KC	Basis	KC	FI	HZ	SM	-
Chlorpromazin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Chlorprothixen	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Chostridium difficile GDH	Stuhl		CLIA	90	3	1	-	-	Spez	-	-	-	-	-
Chostridium difficile Toxin	Stuhl		CLIA	90	3	1	-	-	Spez	-	-	-	-	-
Cholesterol, gesamt	Serum		PHOT	90	7	7	ANA	KC	Basis	KC	FI	HZ	SM	-
Cholinesterase	Serum		PHOT	360	7	6h	ANA	KC	Basis	KC	-	-	-	-
cis-Flupentixol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Citalopram	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-
Clobazam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Clomethiazol	Serum		LC-MS/MS	90	1	5h	-	-	-	HPLC	-	-	-	-
Clonazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Clozapin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
CMV-IgG i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-
CMV-IgG-Ak	Serum		ECLIA	180	28	7	-	-	Basis	-	-	-	-	-
CMV-IgG-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-
CMV-IgM-Ak	Serum		ECLIA	90	28	7	-	-	Basis	-	-	-	-	-
CMV-IgM-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-
cN-1A	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-
Coeruloplasmin	Serum		NEPH	90	7	-	-	-	Liq	-	-	-	-	-
CO-Hämoglobin	EDTA		PHOT	>120	120	3	-	-	-	Häma	-	-	-	-
Coronaviren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-
Coronavirus SARS-CoV-2 (PCR)	Abstrich		PCR	-	7	2	-	-	PCR	Häma	FI	HZ	-	-
Coronavirus SARS-CoV-2 IgG	Serum		CLIA	>5	5	1	-	-	Spez	-	-	-	-	-
Corticotropin (ACTH)°	EDTA-P		ECL	70	3h	2h	-	KC	-	-	-	-	-	-
Cortisol i.S.	Serum		ECL	360	4	1	-	KC	Basis	KC	-	-	-	-
Coxsackievirus IgA	Serum		ELISA	>7	7	-	-	Sero	-	-	-	-	-	-
Coxsackievirus IgG	Serum		ELISA	>7	7	-	-	Sero	-	-	-	-	-	-
Coxsackievirus IgM	Serum		ELISA	>7	7	-	-	Sero	-	-	-	-	-	-

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich									
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM			
Bezeichnung																
C-Peptid	Serum		ECL	30	1	4h	-	KC	Basis	KC	-	-	-	-	-	-
C-reaktives Prot. (hoch sens.)	Serum		TURB	>3a	60	11	-	-	Basis	-	-	-	-	-	-	-
C-reaktives Protein quant.	Serum		TURB	360	21	14	ANA	KC	Basis	KC	FI	HZ	SM			
Creatin-Kinase (CK)	Serum		PHOT	28	7	2	ANA	KC	Basis	KC	FI	HZ	SM			
Creatin-Kinase-MB (CK-MB)	Serum		PHOT	28	8	8h	ANA	KC	Basis	KC	FI	HZ	SM			
Cryptosporidium spp. (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-	-
Cyclospora cayetanensis (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-	-
Cyclosporin A	EDTA		LC-MS/MS	28	7	1	-	-	-	HPLC	-	-	-	-	-	-
CYFRA 21-1	Serum		ECL	180	30	7	-	-	Basis	-	-	-	-	-	-	-
Cystatin C i.S.	Serum		TURB	2a	7	7	ANA	KC	Basis	KC	-	-	-	-	-	-
Cytomegalievirus-DNA (PCR)	PRÜFEN!		PCR	90	14	7	-	-	PCR	-	-	-	-	-	-	-
D-Dimere	Citrat		TURB	180	4	8h	ANA	KC	Basis	Häma	FI	HZ	-	-	-	-
Dehydroripirozol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Demoxepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Desalkylflurazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Desethylamiodaron	Serum		HPLC	28	4	-	-	-	-	HPLC	-	-	-	-	-	-
Desmethylclobazam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Desmethylflunitrazepam	Serum		LC/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Desmethylmesuximid	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Desmethylmianserin	Serum		LC/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-	-
Desmethylmirtazapin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-	-
DHEA-S	Serum		ECL	360	14	5	-	-	Basis	-	-	-	-	-	-	-
Diazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Dientamoeba fragilis (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-	-
Differenzialblutbild	EDTA	nach Abnahme: 10 h		-	-	10h	ANA	Häma	Basis	Häma	FI	HZ	SM			
Digitoxin	Serum		ECL	180	14	4	-	-	Basis	KC	-	-	-	-	-	-
Digoxin	Serum		KIMS	14	1	-	-	-	-	KC	-	-	-	-	-	-
Dihydropropion	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-	-
Dihydrotestosteron (DHT)	Serum		ELISA	>1	1	4h	-	Sero	-	-	-	-	-	-	-	-
Diphtherie-Toxoid: AK (IgG)	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-	-	-	-
Dir. Coombs Test C3c-spezifisch	Vollblut		Aggl	-	10	1	-	-	-	Häma	-	-	-	-	-	-
Dir. Coombs Test C3d-spezifisch	Vollblut		Aggl	-	10	1	-	-	-	Häma	-	-	-	-	-	-
Dir. Coombs Test IgA-spezifisch	Vollblut		Aggl	-	10	1	-	-	-	Häma	-	-	-	-	-	-
Dir. Coombs Test IgG-spezifisch	Vollblut		Aggl	-	10	1	-	-	-	Häma	-	-	-	-	-	-
Dir. Coombs Test IgM-spezifisch	Vollblut		Aggl	-	10	1	-	-	-	Häma	-	-	-	-	-	-
Direkter Coombstest polysp.	Vollblut		Aggl	-	10	1	-	-	-	Häma	FI	HZ	-	-	-	-
Dosulepin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-	-
dRVVT (dilute Russel viper venom time)	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-	-	-	-
Duloxetine	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-	-
E. coli O157 (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-	-
Early Antigen (EA)-IgG-AK	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-	-	-	-
EBV(VCA/EA)-IgM-AK	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-	-	-	-
EBV: IgG-Blot	Serum		IB	>14	14	1	-	Sero	-	-	-	-	-	-	-	-
EBV-IgG-Ak i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-	-
EBV-IgG-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-	-
EBV-VCA-IgM-Ak i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-	-
EBV-VCA-IgM-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-	-
Echovirus IgA	Serum		ELISA	>7	7	-	-	Sero	-	-	-	-	-	-	-	-
Echovirus IgG	Serum		ELISA	>7	7	-	-	Sero	-	-	-	-	-	-	-	-
Echovirus IgM	Serum		ELISA	>7	7	-	-	Sero	-	-	-	-	-	-	-	-
ECP	Serum		EIA	>5	5	2	-	-	-	Allerg	-	-	-	-	-	-
Ecstasy i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-	-	-	-
Eisen	Serum		PHOT	>2a	21	7	ANA	KC	Basis	KC	FI	HZ	SM			
Eiweiß i.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM			
EJ	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-	-
Entamoeba histolytica (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-	-
Entero-/Rhinoviren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-	-	-
Enterovirus-RNA (Cox,Echo,Polio)	Stuhl		PCR	-	5	2	-	-	PCR	-	-	-	-	-	-	-
Eosinophile, absolut	EDTA		WhF	-	2	2	ANA	Häma	Basis	Häma	FI	HZ	-	-	-	-
Epstein-Barr-Virus-AK	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-	-	-	-
Epstein-Barr-Virus-DNA (PCR)	EDTA		PCR	90	14	7	-	-	PCR	-	-	-	-	-	-	-
Erythropoetin	Serum		ELISA	>1	1	8h	-	KC	-	-	-	-	-	-	-	-
Erythrozyten	EDTA		WhF	-	3	3	ANA	Häma	Basis	Häma	FI	HZ	SM			
Erythrozyten i.H. (mikrosk.)	Urin		Mikr/Elz	-	1	4h	-	-	-	-	FI	HZ	SM			
Erythrozyten i.H. (Zellzählg.)	Urin		Mikr/Elz	-	1	4h	ANA	Häma	Basis	KC	-	-	-	-	-	-
Erythrozyten/Hämoglobin i.H.	Urin		Mikr/Elz	-	1	4h	ANA	Häma	Basis	KC	FI	HZ	SM			
Ery-Verteilungsbreite (RDW)	EDTA		RECH	-	1	1	ANA	Häma	Basis	Häma	FI	HZ	-	-	-	-
Estazolam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Ethosuximid	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Everolimus	EDTA		LC-MS/MS	28	7	1	-	-	-	HPLC	-	-	-	-	-	-
Faktor V Leiden-Mutation (G1691A)	EDTA		PCR	-	14	7	-	-	PCR	-	-	-	-	-	-	-
Felbamat	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Ferritin	Serum		ECL	360	7	2	ANA	KC	Basis	KC	-	-	-	-	-	-
Fibrinogen	Citrat		KOAG	540	-	4h	ANA	KC	-	Häma	FI	HZ	-	-	-	-
Flunitrazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Fluoxetin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-	-
Fluphenazin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Flurazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-	-
Fluvoxamin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-	-
Follitropin (FSH)	Serum		ECL	180	14	5	-	KC	Basis	-	-	-	-	-	-	-

Messgröße Bezeichnung	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich						
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM
Folsäure	Serum	lichtgeschützt	ECL	28	2	2h	ANA	KC	Basis	KC	-	-	-
freie Kappa-Leichtketten i.S.	Serum		NEPH	90	21		-	-	Liq	-	-	-	-
freie Lambda-Leichtketten i.S.	Serum		NEPH	90	21		-	-	Liq	-	-	-	-
Fruktose Intoleranz	EDTA		PCR	-	>7	7	-	PCR	-	-	-	-	-
FSME-IgG i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
FSME-IgG-Ak	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
FSME-IgG-AK-Index	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-
FSME-IgM i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
FSME-IgM-Ak	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
FSME-IgM-AK-Index	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-
Gabapentin i.S.	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Gamma-Globulin (%)	Serum		ELPHO	60	10	3	-	Sero	Basis	-	-	-	-
Gamma-GT	Serum		PHOT	360	7	7	ANA	KC	Basis	KC	FI	HZ	SM
Gentamicin	Serum		KIMS	28	7	1	-	-	-	KC	-	-	-
Ger.-Faktor II	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor IX	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor V	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor VII	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor VIII (Aktivität)*	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor X	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor XI	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor XII	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ger.-Faktor XIII (Antigen)	Citrat		KOAG	30	12h	4h	-	KC	-	-	-	-	-
Ges./β-HCG	Serum		ECL	360	14	5	-	KC	Basis	KC	-	HZ	-
Gesamtprotein	Liquor		PHOT	360	6	1	-	-	Basis	-	-	-	-
Gewebstransglutaminase IgA-Ak	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-
Gewebstransglutaminase IgG-Ak	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-
Giardia lamblia (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-
GLDH	Serum		PHOT	28	7	7	-	KC	-	-	-	-	-
Glucose i.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM
Glucose i.H.	Urin		PHOT	-	1	4h	-	-	Basis	KC	-	-	-
Glukose i. Pl. (NaF)	Fluorid		PHOT	-	7	3	-	KC	Basis	Häma	-	-	SM
Glukose i.S.	Serum		PHOT	-	3	8h	ANA	KC	-	KC	FI	HZ	-
Glukose, kapillär (hämol.)	Kapillarblut		AMPE	-	-	30'	ANA	-	Basis	-	-	-	-
Glukose, POCT	Kapillarblut		AMPE	-	-	-	-	-	-	Häma	FI	HZ	-
Großes Blutbild	EDTA	nach Abnahme: 10 h		-	-	10h	ANA	Häma	Basis	Häma	FI	HZ	SM
Guanfacin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Haloperidol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Hämatokrit	EDTA		WhF	-	1	1	ANA	Häma	Basis	Häma	FI	HZ	SM
Hämochromatose (HFE)	EDTA		PCR	-	14	7	-	-	PCR	-	-	-	-
Hämoglobin	EDTA		SLS	-	3	3	ANA	Häma	Basis	Häma	FI	HZ	SM
Hämoglobin im Stuhl	Stuhl		Immunoass	14	14	5	-	-	Spez	-	-	-	-
Haptoglobin	Serum		NEPH	90	7		-	-	Liq	-	-	-	-
Harnanalyse (semiquantitativ)	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM
Harnsäure i.H.	Urin		PHOT	>4	4	1	-	-	-	KC	-	-	-
Harnsäure i.S.	Serum		PHOT	180	7	3	ANA	KC	Basis	KC	FI	HZ	SM
Harnstoff i.H.	Urin		PHOT	30	7	2	-	-	Basis	KC	-	-	-
Harnstoff i.S.	Serum		PHOT	360	7	7	ANA	KC	Basis	KC	FI	HZ	SM
HbA1c (gemäß IFCC)	EDTA		Elpho	90	7	3	-	Häma	Basis	Häma	-	-	-
HBe-Antigen	Serum		ECLIA	90	11	7	-	-	Basis	-	-	-	-
HBs-Ag Bestätigungstest	Serum		NT	180	14	7	-	-	Basis	-	-	-	-
HBs-Antigen	Serum		ECLIA	180	14	7	ANA	KC	Basis	KC	-	-	-
HDL-Cholesterol	Serum		PHOT	360	7	3	ANA	KC	Basis	KC	FI	HZ	-
HE4 (Hum. Epididymis Prot. 4) i.S.	Serum		ECLIA	84	2	5h	-	-	Basis	KC	-	-	-
Helicobacter pylori Ak IgA	Serum		ELISA	>14	14	4h	-	-	-	Viro	-	-	-
Helicobacter pylori Ak IgG	Serum		ELISA	>14	14	4h	-	-	-	Viro	-	-	-
Helicobacter-Blot-IgA°	Serum		IB	>14	14	4h	-	-	-	Viro	-	-	-
Helicobacter-Blot-IgG°	Serum		IB	>14	14	4h	-	-	-	Viro	-	-	-
Heparin/PF-4-Komplex-AK (HIT)	Serum		IC	>90	30	1	-	-	-	Häma	-	-	-
Hepatitis-B-Virus DNA, quant. (Viruslast)	EDTA		PCR	-	14	2	-	-	PCR	-	-	-	-
Hepatitis-C-Virus RNA, quant. (Viruslast)	EDTA		PCR	-	14	2	-	-	PCR	-	-	-	-
Hepatitis-C-Virus: Antikörper (Bestätigungstest)	Serum		IB	>14	14	1	-	-	-	Viro	-	-	-
Hepatitis-C-Virus-Ak	Serum		ECLIA	90	14	7	ANA	KC	Basis	KC	-	-	-
Hepatitis-E-IgG-AK	Serum		CLIA	>7	7	1	-	-	Spez	-	-	-	-
Hepatitis-E-IgM-AK	Serum		CLIA	>7	7	1	-	-	Spez	-	-	-	-
HFR (Unreife Retikulozyten)	EDTA		FC	-	-	1	-	Häma	-	-	-	-	-
HIV-1/2 AK Western Blot	Serum		IB	>14	14	1	-	-	-	Viro	-	-	-
HIV-1-RNA (ultrasensitiv, PCR)	EDTA PCR		PCR	-	14	2	-	-	PCR	-	-	-	-
HLA-B27 (PCR)	EDTA		PCR	30	14	7	-	-	PCR	-	-	-	-
HLA-DQ2/DQ8 (Zöliakie)	EDTA		PCR	-	>7	7	-	PCR	-	-	-	-	-
HLA-DRB1 PCR	EDTA		PCR	-	>7	7	-	PCR	-	-	-	-	-
Holotranscobalamin	Serum		ELISA	70	3	16h	-	KC	-	KC	-	-	-
Homocystein	HCY-R.		LC-MS/MS	90	7	2	-	-	-	HPLC	-	-	-
HPV-High Risk Genotyp°	Abstrich		TMA	-	90	30	-	-	PCR	-	-	-	-
HPV-mRNA (High Risk)°	Abstrich		TMA	-	90	30	-	-	PCR	-	-	-	-
HSV-1/2-IgA i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
HSV-1/2-IgA-(Screen)-AK-Index	Liquor		ELISA				-	-	Liq	-	-	-	-
HSV-1/2-IgG i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
HSV-1/2-IgG-(Screen)-AK-Index	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-
HSV-1/2-IgM i.S.	Serum		ELISA	90	7		-	-	-	-	-	-	-

Messgröße Bezeichnung	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich							
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM	
HSV-1/2-IgM-(Screen)-AK-Index	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-	-
HSV-1/2-IgM-Ak	Serum		CLIA	90	7		-	-	Spez	-	-	-	-	-
HSV-1-IgG-Ak	Serum		CLIA	90	7		-	-	Liq	-	-	-	-	-
HSV-2-IgG-Ak	Serum		CLIA	90	7		-	-	Liq	-	-	-	-	-
HSV-I-DNA (PCR)	Abstrich		PCR	90	14	7	-	-	PCR	-	-	-	-	-
HSV-II-DNA (PCR)	Abstrich		PCR	90	14	7	-	-	PCR	-	-	-	-	-
Hydroxybupropion	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-
IgA im L.c.	Liquor		TURB	180	7		-	-	Liq	-	-	-	-	-
IgA-Rheumafaktor i. S.	Serum		EIA	>5	5	1	-	Sero	-	-	-	-	-	-
IGF-BP3 (IGF-binding Prot. 3)°	Serum		CLIA	>1	-	1h	-	Sero	-	-	-	-	-	-
IGF-I (Somatomedin C)°	Serum		CLIA	>1	-	1h	-	Sero	-	-	-	-	-	-
IgG im L.c.	Liquor		TURB	180	7		-	-	Liq	-	-	-	-	-
IgG-Rheumafaktor i. S.	Serum		EIA	>5	5	1	-	Sero	-	-	-	-	-	-
IgM im L.c.	Liquor		TURB	180	7		-	-	Liq	-	-	-	-	-
IgM-Rheumafaktor i. S.	Serum		EIA	>5	5	1	-	Sero	-	-	-	-	-	-
Immundefixation i.S.	Serum		ELPHO	>14	14	3	-	-	Spez	-	-	-	-	-
Immunglobulin A	Serum		TURB	240	240	240	-	KC	Basis	KC	-	-	-	-
Immunglobulin G	Serum		TURB	240	240	120	-	KC	Basis	KC	-	-	-	-
Immunglobulin G i.H.	Urin		NEPH	-	-	1	-	-	Liq	-	-	-	-	-
Immunglobulin M	Serum		TURB	180	120	60	-	KC	Basis	KC	-	-	-	-
Immunglobulin-E	Serum		EIA	>7	7	2	-	-	-	Allerg	-	-	-	-
Influenza A (saisonal)-RNA (PCR)	Abstrich		PCR	-	7	2	-	-	PCR	Häma	FI	HZ	-	-
Influenza A-Viren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-
Influenza B-RNA (PCR)	Abstrich		PCR	-	7	2	-	-	PCR	Häma	FI	HZ	-	-
Influenza B-Viren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-
INR	Citrat		KOAG	180	-	1	ANA	KC	Basis	Häma	FI	HZ	SM	-
Insulin i.S.	Serum		ECL	180	2	4h	-	KC	-	-	-	-	-	-
Interleukin 6	Serum		ECL	2a	2	6h	-	-	-	KC	-	-	-	-
Interleukin 6 (screen)	Serum		FC	2a	2	8h	-	-	-	-	-	HZ	-	-
JC-Virus-(Polyoma) DNA (PCR) i.S.	Serum		PCR	90	14	7	-	-	PCR	-	-	-	-	-
JC-Virus-(Polyoma) DNA (PCR) im L.c.	Liquor		PCR	90	14	7	-	-	PCR	-	-	-	-	-
Kalium i.H.	Urin		ISE	>14	>14	14	-	-	Basis	KC	-	-	-	-
Kalium i.S.	Serum		ISE	>14	14	14	ANA	KC	Basis	KC	FI	HZ	SM	-
Kell	Vollblut		Aggl	-	10	1	-	-	-	Häma	FI	HZ	-	-
Ketonkörper i.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM	-
Kleines Blutbild	EDTA			-	2	2	ANA	Häma	Basis	Häma	FI	HZ	SM	-
Knochen AP (BAP, Ostase)	Serum		EIA	39m	90	2	-	Sero	-	-	-	-	-	-
Kokain i.H.(immunol.)	Urin		KIMS	>5	5	1	-	-	-	KC	-	-	-	-
Kokain i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-	-
Kreatinin i.H.	Urin		PHOT	180	6	2	-	KC	Basis	KC	-	-	-	-
Kreatinin i.S.	Serum		Jaffe	90	7	7	ANA	KC	Basis	KC	FI	HZ	SM	-
Kupfer i.S.	Serum		PHOT	360	14	1	-	KC	-	-	-	-	-	-
Lacosamid	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Lactat	Fluorid	sofort ins Labor, zentrifug.	PHOT	38	14	8h	-	-	Basis	-	FI	-	-	-
Lactat im L.c.	Liquor	zellarmer L.c.: 24 h	PHOT	38	14	3h	-	-	-	-	-	-	-	-
Lactose Intoleranz	EDTA		PCR	-	7	-	-	PCR	-	-	-	-	-	-
Lamotrigin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
LDH	Serum		PHOT	42	4	7	ANA	KC	Basis	KC	FI	HZ	SM	-
LDL-Cholesterol	Serum		PHOT	360	7	1	ANA	KC	Basis	KC	FI	HZ	-	-
Leber-Ak	Serum		IFT	>14	14	1	-	-	-	Viro	-	-	-	-
Legionella pneum. IgG (Gr.1)°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-	-
Legionella pneum. IgM (Gr.1)°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-	-
Legionella pneumophila Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-
Leukozyten	Liquor		MIKR	-	-	2h	-	-	Liq	-	-	-	-	-
Leukozyten	EDTA		WhF	-	3	3	ANA	Häma	Basis	Häma	FI	HZ	SM	-
Leukozyten i.H. (mikrosk.)	Urin		Mikr/Elz	-	1	4h	-	-	-	-	FI	HZ	SM	-
Leukozyten i.H. (Zellzählg.)	Urin		Mikr/Elz	-	1	4h	ANA	KC	Basis	KC	-	-	-	-
Leukozyten/Esterase i.H.	Urin		Mikr/Elz	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM	-
Leukozytenkonglomerate i.H.	Urin		Mikr/Elz	-	1	4h	ANA	-	-	-	-	-	-	-
Levetiracetam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Levomepromazin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
LFR (Reife Retikulozyten)	EDTA		FC	-	-	1	-	Häma	-	-	-	-	-	-
Lipase	Serum		PHOT	360	7	7	ANA	KC	Basis	KC	FI	HZ	SM	-
Lipoprotein (a)	Serum		TURB	>2	2	8h	-	KC	Basis	KC	-	-	-	-
Liquorzytologie	Liquor		MIKR	-	-	2h	-	-	Liq	-	-	-	-	-
Lithium	Serum		PHOT	>7	7	1	-	-	Basis	KC	-	-	-	-
Lorazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Lormetazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-
Lutropin (LH)	Serum		ECL	180	14	5	-	KC	Basis	-	-	-	-	-
Lymphozyten, absolut	EDTA		WhF	-	2	2	ANA	Häma	Basis	Häma	FI	HZ	-	-
Lymphozyten, absolut	EDTA		WhF/SCG	-	2	2	-	-	-	-	-	-	SM	-
Magnesium	Serum		PHOT	360	7	7	-	KC	Basis	KC	FI	-	-	-
Magnesium i.H.	Urin		PHOT	360	3	3	-	-	Basis	-	-	-	-	-
Masern-Ak-IgG	Serum		ELISA	90	7		-	-	Liq	-	-	-	-	-
Masern-Ak-IgM	Serum		ELISA	90	7		-	-	Liq	-	-	-	-	-
Masern-IgG i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-	-
Masern-IgG-AK-Index	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-	-
Masern-IgM i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-	-
Masern-IgM-AK-Index	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-	-
MDA5	Serum		BLOT	>14	14		-	-	-	Viro	-	-	-	-

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich						
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM
Bezeichnung													
Medazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Melperon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Metanephrin i. Pl. (LC/MS)	EDTA-P		LC/MS	28	3	-	-	-	-	HPLC	-	-	-
Metapneumoviren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-
Methadon i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-
Methämoglobin	EDTA		PHOT	30	2	1h	-	-	-	Häma	-	-	-
Methamphetamine i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-
Methoxytyramin i. Pl. (LC/MS)	EDTA-P		LC/MS	28	3	-	-	-	-	HPLC	-	-	-
Methylmalonsäure	Serum		LC/MS	360	180	7	-	-	-	HPLC	-	-	-
Methylphenidat	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
MFR (Mittlereife Retikulozyten)	EDTA		FC	-	-	1	-	Häma	-	-	-	-	-
Mi-2α	Serum		BLOT	>14	14		-	-	-	Viro	-	-	-
Mi-2β	Serum		BLOT	>14	14		-	-	-	Viro	-	-	-
Mianserin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Midazolam	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Milnacipran	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Mirtazapin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Mittl. Erythrozytenvolumen (MCV)	EDTA		WhF	-	1	1	ANA	Häma	Basis	Häma	FI	HZ	SM
Mittl. korp. Hämoglobin (MCH)	EDTA		RECH	-	3	3	ANA	Häma	Basis	Häma	FI	HZ	-
Mittl. korp. Hämoglobinkonz. (MCHC)	EDTA		RECH	-	3	3	ANA	Häma	Basis	Häma	FI	HZ	-
Moclobemid	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Mono / Baso / Eos (abs.)	EDTA		WhF/SCG	-	2	2	-	-	-	-	-	-	SM
Monozyten, absolut	EDTA		WhF	-	2	2	ANA	Häma	Basis	Häma	FI	HZ	-
MRSA	Abstrich		PCR	-	7	1	-	-	Mibi	-	-	-	-
MTHFR (C677T)	EDTA		PCR	-	>7	7	-	PCR	-	-	-	-	-
Mumps-IgG i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
Mumps-IgG-Ak	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
Mumps-IgG-AK-Index	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-
Mumps-IgM i.S.	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
Mumps-IgM-Ak	Serum		ELISA	90	7		-	-	Liq	-	-	-	-
Mumps-IgM-AK-Index°	Liquor		ELISA	90	7		-	-	Liq	-	-	-	-
Mycoplasma genitalium Multiplex	Abstrich		PCR	-	7	2	-	-	Mibi	-	-	-	-
Mycoplasma hominis Multiplex	Abstrich		PCR	-	7	2	-	-	Mibi	-	-	-	-
Mycoplasma pneumoniae Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-
Mykoplasmen-Ak-IgA	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-
Mykoplasmen-Ak-IgG	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-
Mykoplasmen-Ak-IgM	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-
Myoglobin	Serum		ECL	360	14	8	-	KC	Basis	KC	FI	HZ	-
Natrium i.H.	Urin		ISE	>14	>14	14	-	-	Basis	KC	-	-	-
Natrium i.S.	Serum		ISE	>14	14	14	ANA	KC	Basis	KC	FI	HZ	SM
N-Desalkylquetiapin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
N-Desmethylcitalopram	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
N-Desmethylolanzapin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
N-Desmethylsertralin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Nefazodon	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Neisseria gonorrhoeae Multiplex	Abstrich		PCR	-	7	2	-	-	Mibi	-	-	-	-
Neisseria gonorrhoeae rRNA	Abstrich		NAT	-	-	30	-	-	PCR	-	-	-	-
Neisseria gonorrhoeae rRNA	Urin		NAT	-	-	30	-	-	PCR	-	-	-	-
Neutrophile, absolut	EDTA		WhF	-	2	2	ANA	KC	Basis	Häma	FI	HZ	-
Neutrophile, absolut	EDTA		WhF/SCG	-	2	2	-	-	-	-	-	-	SM
Nitrazepam	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Nitrit i.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM
NK-Lymph. (CD56)	EDTA		FC	-	-	1	-	-	Spez	-	-	-	-
Norclozapin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Nordiazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Norfluoxetin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Normetanephrin i.Pl. (LC/MS)	EDTA-P		LC/MS	28	3	-	-	-	-	HPLC	-	-	-
Norovirus Genogruppe I (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-
Norovirus Genogruppe II (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-
Norovirus Genotyp 2	Stuhl		PCR	-	5	2	-	-	PCR	-	-	-	-
Norovirus Genotyp1	Stuhl		PCR	-	5	2	-	-	PCR	-	-	-	-
NSE	Serum		ECLIA	90	5	2	-	-	-	KC	-	-	-
NSE im L.c.°	Liquor		ECLIA	90	5	2	-	-	-	KC	-	-	-
NT-proBNP	Serum		ECL	2a	6	3	ANA	KC	Basis	KC	-	-	-
Nukl. Antigen (EBNA1)-IgG-AK	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-
NXP2	Serum		BLOT	>14	14		-	-	-	Viro	-	-	-
O-Desmethyltramadol	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
O-Desmethylvenlafaxin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
OJ	Serum		BLOT	>14	14		-	-	-	Viro	-	-	-
Olanzapin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Oligoklonale Banden (IEF)	Liquor		IEF	90	14		-	-	Liq	-	-	-	-
Opiate i.H.(immunol.)	Urin		KIMS	>5	5	1	-	-	-	KC	-	-	-
Opiate i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-
Opiamol	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Osmolalität i.H.°	Urin		OSMO	-	-	1	-	-	-	-	-	-	SM
Osmolalität i.S.°	Serum		OSMO	-	-	1	-	-	-	-	-	-	SM
Osteocalcin	Serum		CLIA	>1	-	1	-	Sero	-	-	-	-	-
Oxazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Oxcarbazepin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
p - ANCA (IFT)	Serum		IFT	>14	14	-	-	-	-	Viro	-	-	-

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich								
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM		
Bezeichnung															
P-ANCA (MPO)	Serum		ELISA	-	14	7	-	Sero	Spez	-	-	-	-	-	-
Pankreas-Amylase	Serum		PHOT	>30	30	7	ANA	KC	Basis	KC	FI	HZ	SM		
Pankreas-Elastase i. Stuhl	Stuhl		CLIA	360	7	5	-	-	Mibi	-	-	-	-	-	-
Parainfluenzaviren Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-	-
Parathormon	Serum		ECL	180	2	8h	-	KC	Basis	KC	-	-	-	-	-
Parathormon im EDTA-Plasma	EDTA		ECL	180	3	2	-	-	-	KC	-	-	-	-	-
Parietalzell-AK	Serum		IB	>5	5	-	-	Sero	-	-	-	-	-	-	-
Paroxetin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-
Parvovirus B19 IgG-Ak	Serum		CLIA	>7	7	1	-	-	Spez	-	-	-	-	-	-
Parvovirus B19 IgM-Ak	Serum		CLIA	>7	7	1	-	-	Spez	-	-	-	-	-	-
Parvovirus-B19-IgG Blot	Serum		IB	>7	7	1	-	-	-	-	-	-	-	-	-
Parvovirus-B19-IgM Blot	Serum		IB	>7	7	1	-	-	-	-	-	-	-	-	-
PCO 2	Kapillarblut		AMPE	-	-	10'	-	-	-	Häma	FI	HZ	-	-	-
pCO2-Wert	Kapillarblut		ELEK	-	-	10'	-	-	Basis	-	-	-	-	-	-
Perampanel	Serum		LC/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Perazin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Phenobarbital	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Phenytoin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Phosphat i.H.	Urin		PHOT	>180	180	1	-	-	Basis	KC	-	-	-	-	-
Phosphat i.S.	Serum		PHOT	360	4	1	ANA	KC	Basis	KC	-	-	-	SM	
Phospho-Tau (181P) i.L.c.	Liquor		CMIA	60	7	-	-	-	Liq	-	-	-	-	-	-
pH-Wert	Kapillarblut		ELEK	-	-	10'	-	-	Basis	-	-	-	-	-	-
pH-Wert i.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM		
Pipamperon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
PL-12	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
PL-7	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
PM-Scl100	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
PM-Scl75	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
PO 2	Kapillarblut		AMPE	-	-	10'	-	-	-	Häma	FI	HZ	-	-	-
pO2-Wert	Kapillarblut		ELEK	-	-	10'	-	-	Basis	-	-	-	-	-	-
Prazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Pregabalin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Primidon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Procalcitonin	Serum		ELFA	390	2	1	-	KC	Basis	KC	FI	HZ	-	-	-
Progesteron	Serum		ECL	180	5	1	-	KC	Basis	-	-	-	-	-	-
Prokollagen-I-N-term. Propeptid	Serum		ECLIA	180	5	1	-	-	Basis	-	-	-	-	-	-
Prolaktin	Serum		ECL	180	14	5	-	KC	Basis	-	-	-	-	-	-
Promethazin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Prostata-spez. AG (PSA)	Serum		ECL	168	5	1	ANA	KC	Basis	KC	-	-	-	-	-
Prostata-spez. AG, frei (fPSA)	Serum		ECL	84	5	8h	-	KC	Basis	KC	-	-	-	-	-
Protein 14-3-3 im L.c. (ELISA)	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Protein C (Akt.)	Citrat		CHRO	30	12h	4h	-	KC	-	-	-	-	-	-	-
Protein i.H.	Urin		TURB	30	7	1	-	KC	Basis	KC	-	-	-	-	-
Protein S, frei (Antigen)	Citrat		TURB	30	12h	4h	-	KC	-	-	-	-	-	-	-
Protein S-100 i.S.	Serum		ECL	90	2	8h	-	-	Basis	-	-	-	-	-	-
Protein, gesamt, i.S.	Serum		PHOT	360	28	6	ANA	KC	Basis	KC	FI	HZ	SM		
Prothipendyl	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Prothrombin-Mutation (G20210A)	EDTA		PCR	-	14	7	-	-	PCR	-	-	-	-	-	-
PTT	Citrat		KOAG	180	-	4h	ANA	KC	Basis	Häma	FI	HZ	SM		
PTT, lupusinsensitiv	Citrat		KOAG	180	-	4h	-	KC	-	-	-	-	-	-	-
PTT, lupussensitiv	Citrat		KOAG	180	-	4h	-	KC	-	-	-	-	-	-	-
Quetiapin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Quicktest	Citrat		KOAG	180	-	1	ANA	KC	Basis	Häma	FI	HZ	SM		
R. conori IgG°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-	-	-
R. conori IgM°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-	-	-
Reboxetin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-
Renin	EDTA-P		CLIA	90	-	1	-	-	Spez	-	-	-	-	-	-
Ret-He (Reticul. hemoglobine equiv.)	EDTA		PHOT	-	1	1	-	Häma	Basis	Häma	-	-	-	-	-
Retigabin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Retikulozyten	EDTA		FC	-	1	1	-	Häma	Basis	Häma	-	-	-	-	-
Rhesus-Faktor D	Vollblut		Aggl	-	10	1	-	-	-	Häma	FI	HZ	-	-	-
Rheumafaktor quant.	Serum		TURB	90	8	1	-	-	Basis	KC	-	-	-	-	-
Rh-Formel	Vollblut		Aggl	-	10	1	-	-	-	Häma	FI	HZ	-	-	-
Risperidon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Ritalinsäure	Serum		LC/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-
Ro-52	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
Rotavirus A (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-
Röteln Ak IgG (EIA)	Serum		ECLIA	90	21	7	-	KC	Basis	KC	-	-	-	-	-
Röteln Ak IgM (EIA)	Serum		ECLIA	90	21	3	-	KC	Basis	-	-	-	-	-	-
Röteln-IgG i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Röteln-IgG-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
RSV Multiplex	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	-	-	-
RS-Virus-RNA (PCR)	Abstrich		PCR	-	7	2	-	-	PCR	-	-	-	HZ	-	-
Rufinamid	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
SAE1	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
Salmonella spp. (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-
Sapovirus (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-
SARS-CoV-2 AK IgG Bestätigung	Serum		ELISA	>5	5	1	-	-	-	Viro	-	-	-	-	-
Sauerstoffsättigung	Kapillarblut		AMPE	-	-	10'	-	-	-	Häma	-	-	-	-	-
Saure Phosphatase	Serum		PHOT	120	8	8	ANA	KC	-	-	-	-	-	-	-

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]				Labor - Bereich							
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM		
Bezeichnung															
SCC-Antigen	Serum		ECL	84	14	5	-	-	-	Basis	-	-	-	-	-
Selen i.S.	Serum		AAS	-	30	-	-	AAS	-	-	-	-	-	-	-
Sertindol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Sertralin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-
SHBG	Serum		ECL	360	7	5	-	KC	-	-	-	-	-	-	-
Shiga like toxin 1/2 (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-
Shigella spp. / EIEC (PCR):	Stuhl		PCR	-	5	2	-	-	Mibi	-	-	-	-	-	-
Sirolimus i. B.	EDTA		LC/MS	28	7	1	-	-	-	HPLC	-	-	-	-	-
Somatotropin (STH, hGH)°	Serum		CLIA	>1	-	1	-	Sero	-	-	-	-	-	-	-
spez. Gewicht d.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM		
spezifisches Immunglobulin-E	Serum		EIA	>7	7	2	-	-	-	Allerg	-	-	-	-	-
SRP	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
β2 - Glycoprotein 1 IgG - AK	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-	-	-
β2 - Glycoprotein 1 IgM - AK	Serum		LIA	>7	7	8h	-	Sero	-	-	-	-	-	-	-
β2-Mikroglobulin	Serum		CLIA	90	7	1	-	-	Spez	-	-	-	-	-	-
β-Keton i. Blut	Kapillarblut		AMPE	-	-	-	-	-	-	-	FI	-	-	-	-
β-Trace-Protein	Sekret		NEPH	-	7	-	-	-	Liq	-	-	-	-	-	-
Stiripentol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Sulpirid	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Sultiam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
T.pallidum IgG-Blot i.S.	Serum		IB	90	7	-	-	-	Spez	-	-	-	-	-	-
T.pallidum IgM-Blot i.S.	Serum		IB	90	7	-	-	-	Spez	-	-	-	-	-	-
Tacrolimus II	EDTA		LC-MS/MS	28	7	1	-	-	-	HPLC	-	-	-	-	-
Tau-Protein im L.c.	Liquor		CMIA	60	7	-	-	-	Liq	-	-	-	-	-	-
Temazepam	Serum		LC-MS/MS	28	7	1	-	-	-	HPLC	-	-	-	-	-
Testosteron	Serum		ECL	180	14	5	-	KC	Basis	KC	-	-	-	-	-
Tetanus-Toxoid-Ak	Serum		ELISA	>7	7	1	-	-	-	Viro	-	-	-	-	-
Tetrazepam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
T-Helferz. (CD4)	EDTA		FC	-	-	1	-	-	Spez	-	-	-	-	-	-
T-Helferz. (CD4) absolut	EDTA		FC	-	-	1	-	-	Spez	-	-	-	-	-	-
Theophyllin°	Serum		KIMS	60	7	8h	-	-	-	KC	-	-	-	-	-
Thioridazin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Thrombinzeit	Citrat		KOAG	90	-	4h	ANA	KC	-	Häma	-	-	-	-	-
Thrombozyten	EDTA		WhF	-	2	2	ANA	Häma	Basis	Häma	FI	HZ	SM		
Thymidin-Kinase	Serum		CLIA	90	2	1	-	-	Spez	-	-	-	-	-	-
Thyreoglobulin	Serum		CLIA	720	14	14	-	-	Basis	-	-	-	-	-	-
Thyroxin, frei (fT4)	Serum		ECL	30	7	5	ANA	KC	Basis	KC	FI	HZ	-	-	-
Tiagabin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Tianeptin	Serum		LC/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-
TIF1γ	Serum		BLOT	>14	14	-	-	-	-	Viro	-	-	-	-	-
T-Lymphozyten (CD3)	EDTA		FC	-	-	1	-	-	Spez	-	-	-	-	-	-
Tobramycin°	Serum		PHOT	30	3	-	-	-	-	KC	-	-	-	-	-
Topiramate	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Toxopl.-IgG-Avidität	Serum		LIA	180	7	1	-	-	Spez	-	-	-	-	-	-
Toxoplasmose IgG-Ak	Serum		ECLIA	90	21	3	-	KC	Basis	KC	-	-	-	-	-
Toxoplasmose IgM-Ak	Serum		ECLIA	90	21	3	-	KC	Basis	KC	-	-	-	-	-
Toxoplasmose-IgG i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Toxoplasmose-IgG-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Toxoplasmose-IgM i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Toxoplasmose-IgM-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
TPHA i.S.	Serum		HAH	90	7	-	-	-	Liq	-	-	-	-	-	-
TPHA-Test	Serum		IHA	90	7	-	-	Sero/KC	Liq	-	-	-	-	-	-
Tramadol	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-	-	-
Transferrin	Serum		TURB	180	8	8	ANA	KC	Basis	KC	-	-	-	-	-
Transferrinrezeptor, lösl.	Serum		NEPH	30	7	8h	-	KC	-	-	-	-	-	-	-
Transglutaminase-AK	Serum		IB	>5	5	8h	-	Sero	-	-	-	-	-	-	-
Tranlycypromin	Serum		LC-MS/MS	90	1	5h	-	-	-	HPLC	-	-	-	-	-
Trazodon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Trepon. pall. IgG-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Trepon. pall. IgM-AK-Index	Liquor		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Treponema pall.-IgG i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Treponema pall.-IgG i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Treponema pall.-IgM i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Treponema pall.-IgM i.S.	Serum		ELISA	90	7	-	-	-	Liq	-	-	-	-	-	-
Treponema pallidum-Suchtest	Serum		ECLIA	360	14	7	-	-	-	-	-	-	-	-	-
Triazolam	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-	-	-
Trichomonas Vaginalis Multiplex	Abstrich		PCR	-	7	2	-	-	Mibi	-	-	-	-	-	-
Tricycl. Antidepressiva i.H., qual.	Urin		IC	>5	5	1	-	Sero	-	-	FI	HZ	-	-	-
Triglyceride	Serum		PHOT	90	10	2	ANA	KC	Basis	KC	FI	HZ	SM		
Trijodthyronin, frei (fT3)	Serum		ECL	30	7	5	ANA	KC	Basis	KC	FI	HZ	-	-	-
Troponin T, hochsensitiv	Serum		ECL	360	1	<1	ANA	KC	Basis	KC	FI	HZ	-	-	-
Tryptase	Serum		EIA	>7	7	2	-	-	-	Allerg	-	-	-	-	-
TSH (basal)	Serum		ECL	2a	14	8	ANA	KC	Basis	KC	FI	HZ	-	-	-
T-Spot.COVID°	Hep.-Blut		LTT	-	-	1	-	-	Spez	-	-	-	-	-	-
T-Spot.Tb-Test (M. tuberculosis)°	Hep.-Blut		LTT	-	-	2	-	-	Spez	-	-	-	-	-	-
T-Suppressor (CD8)	EDTA		FC	-	-	1	-	-	Spez	-	-	-	-	-	-
Ureaplasma parvum Multiplex	Abstrich		PCR	-	7	2	-	-	Mibi	-	-	-	-	-	-
Ureaplasma urealyticum (PCR):	Abstrich		PCR	-	7	2	-	-	Mibi	-	-	-	-	-	-
Urobilinogen i.H.	Urin		TS	-	1	4h	ANA	KC	Basis	KC	FI	HZ	SM		
Valproinsäure	Serum		PHOT	90	7	2	-	-	Basis	-	-	-	-	-	-

Messgröße	Material	Besonderheiten	Verfahren	Stabilität [d]			Labor - Bereich						
				TK	KS*	RT	ANA	C	DD	EE	FI	HZ	SM
Valproinsäure	Serum		LC-MS/MS	90	7	2	-	-	-	HPLC	-	-	-
Vancomycin°	Serum		KIMS	360	14	2	-	-	-	KC	-	-	-
Varizella-Zoster-Virus DNA (PCR)	EDTA		PCR	90	14	7	-	-	PCR	-	-	-	-
VDRL (qual.)	Serum		AGGL	90	7		-	-	Liq	-	-	-	-
Venlafaxin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
Virus capsid (VCA) p18-IgG-AK	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-
Virus capsid (VCA) p23-IgG-AK	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-
Vitamin B1 (Thiamin)	EDTA	lichtgeschützt	HPLC	180	1	-	-	-	HPLC	-	-	-	-
Vitamin B12 (Cobalamin)	Serum	lichtgeschützt	ECL	56	2	2h	ANA	-	KC	Basis	KC	-	-
Vitamin B2 (Riboflavin)	EDTA	lichtgeschützt	HPLC	>1	1	-	-	-	HPLC	-	-	-	-
Vitamin B6 (Pyridoxin)	EDTA	lichtgeschützt	HPLC	>60	7	-	-	-	HPLC	-	-	-	-
Vortioxetin	Serum		LC-MS/MS	90	7	1	-	-	-	HPLC	-	-	-
vWF: Aktivität (Willebrand-Faktor-Aktivität)	Citrat		TURB	30	12h	4h	-	-	KC	-	-	-	-
vWF: Antigen (Willebrand-Faktor-Ag)	Citrat		TURB	30	12h	4h	-	-	KC	-	-	-	-
VZV-IgA i.S.	Serum		ELISA	90	7		-	-	-	Liq	-	-	-
VZV-IgA-AK-Index	Liquor		ELISA	90	7		-	-	-	Liq	-	-	-
VZV-IgG i.S.	Serum		ELISA	90	7		-	-	-	Liq	-	-	-
VZV-IgG-AK-Index	Liquor		ELISA	90	7		-	-	-	Liq	-	-	-
VZV-IgM i.S.	Serum		ELISA	90	7		-	-	-	Liq	-	-	-
VZV-IgM-AK-Index°	Liquor		ELISA	90	7		-	-	-	Liq	-	-	-
Yersinia enterocolitica (PCR):	Stuhl		PCR	-	5	2	-	-	-	Mibi	-	-	-
Yersinien IgA-Ak (Blot)	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-
Yersinien IgG-Ak (Blot)	Serum		XMAP	>14	14	1	-	-	-	Viro	-	-	-
Zaleplon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Zika-Virus IgG°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-
Zika-Virus IgM°	Serum		CLIA	>7	7	1	-	-	-	Viro	-	-	-
Zink i.S.	Serum		AAS	-	30	-	-	-	AAS	-	-	-	-
Ziprasidon	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Zolpidem	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Zonisamid	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Zopiclon	Serum		LC-MS/MS	90	3h	1h	-	-	-	HPLC	-	-	-
Zotepin	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-
Zuclopendixol	Serum		LC-MS/MS	90	7	3	-	-	-	HPLC	-	-	-